

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending November 8, 2014

Prepared by the Office of Infectious Disease Epidemiology and Outbreak Response Infectious Disease Bureau – Prevention and Health Promotion Administration Maryland Department of Health and Mental Hygiene

The data presented in this document are provisional and subject to change as additional reports come in.

# **SUMMARY**

During the week ending November 8, 2014, influenza-like illness (ILI) intensity in Maryland was **MINIMAL** and there was **LOCAL** geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers and the proportion reported by Emergency Departments both decreased slightly. The proportion of MRITS respondents experiencing ILI increased. Clinical laboratories saw a rise in the proportion of specimens testing positive for influenza. The DHMH lab had 2 specimens test positive for influenza – both were Type A (H3N2). A total of 4 influenza-associated hospitalizations were reported. No respiratory outbreaks were reported. Nationally, influenza activity is low.


Click here to visit our influenza surveillance web page

ILI Intensity Levels			
<b>✓</b> Minimal			
Low			
Moderate			
High			

Influenza Geographic Spread
No Activity
Sporadic
<b>✓</b> Local
Regional
Widespread

#### **ILINet Sentinel Providers**


Seventeen sentinel providers reported a total of 10,761 visits this week. Of those, 122 (1.1%) were visits for ILI. This is below the Maryland baseline of 2.0%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	47 (39%)	48 (36%)	218 (33%)
Age 5-24	56 (46%)	58 (44%)	288 (44%)
Age 25-49	13 (11%)	21 (16%)	101 (15%)
Age 50-64	4 (3%)	5 (4%)	31 (5%)
Age ≥ 65	2 (2%)	1 (1%)	14 (2%)
Total ILI Visits	122 (100%)	133 (100%)	652 (100%)

### Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 43,208 visits this week through the ESSENCE surveillance system. Of those, 642 (1.5%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	217 (34%)	220 (32%)	1145 (29%)
Age 5-24	222 (35%)	237 (35%)	1379 (36%)
Age 25-49	120 (19%)	152 (22%)	879 (23%)
Age 50-64	61 (10%)	54 (8%)	330 (8%)
Age ≥ 65	22 (3%)	21 (3%)	149 (4%)
Unknown	0 (0%)	0 (0%)	1 (<1%)
Total ILI Visits	642 (100%)	684 (100%)	3883 (100%)

#### Other states' influenza information:

Delaware <a href="http://dhss.delaware.gov/dph/epi/influenzahome.html">http://dhss.delaware.gov/dph/epi/influenzahome.html</a>

District of Columbia <a href="http://doh.dc.gov/service/influenza">http://doh.dc.gov/service/influenza</a>

Pennsylvania <a href="http://www.portal.state.pa.us/portal/server.pt/community/influenza\_(flu)/14161">http://www.portal.state.pa.us/portal/server.pt/community/influenza\_(flu)/14161</a>

 Virginia
 http://www.vdh.state.va.us/Epidemiology/Surveillance/Influenza/


 West Virginia
 http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending November 8, 2014

## Community-based Influenza Surveillance (MRITS and Google Flu Trends)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 587 residents responded to the MRITS survey this week. Of those, 10 (1.7%) reported having ILI and missing a cumulative 24 days of regular daily activities. According to Google Flu Trends, influenza activity is "Low" in Maryland.





# **Clinical Laboratory Influenza Testing**

Thirty-two clinical laboratories reported performing 657 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 24 (3.7%) were positive for influenza. Of those testing positive, 15 (62.5%) were influenza Type A and 9 (37.5%) were influenza Type B. The <u>reliability of RIDTs</u> depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	15 (63%)	5 (24%)	37 (44%)
Туре В	9 (38%)	16 (76%)	47 (56%)
Total	24 (100%)	21 (100%)	84 (100%)



#### **State Laboratories Administration Influenza Testing**

The DHMH Laboratories Administration performed a total of 48 PCR tests for influenza and 2 (4.2%) specimens tested positive. Both were influenza Type A (H3N2). PCR testing is more reliable than RIDT. The DHMH testing identifies subtypes of influenza A, information that is not available from the RIDT results. The table below summarizes results by type and subtype.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	-		-
Type A (H3)	2 (100%)	1 (100%)	8 (67%)
Type A (H3N2v)			
Туре В			4 (33%)
Total	2 (100%)	1 (100%)	12 (100%)

# Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2014-15 influenza season? Go to <a href="http://dhmh.maryland.gov/flumd/SitePages/getvaccinated.aspx">http://dhmh.maryland.gov/flumd/SitePages/getvaccinated.aspx</a> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending November 8, 2014

## **Influenza-associated Hospitalizations**


This week, hospitals in Maryland reported 4 influenza-associated hospitalizations. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g. RIDT or PCR, is considered an "influenza-associated hospitalization" for purposes of influenza surveillance.)



Influenza- Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	1 (25%)	1 (20%)	7 (19%)
Age 5-17			1 (3%)
Age 18-24	1 (25%)		1 (3%)
Age 25-49	-	2 (40%)	5 (14%)
Age 50-64	1 (25%)	1 (20%)	8 (22%)
Age ≥ 65	1 (25%)	1 (20%)	15 (41%)
Total	4 (100%)	5 (100%)	37 (100%)

### **Outbreaks of Respiratory Disease**

There were no respiratory outbreaks reported to DHMH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g. from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza		1	1
Influenza-like Illness		1	-
Pneumonia		-	3 (100%)
Other Respiratory			
Total			3 (100%)

#### National Influenza Surveillance (CDC)

During week 45 (November 2 - 8, 2014), influenza activity was low in the United States.

- Viral Surveillance: Of 9,138 specimens tested and reported by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories during week 45, 678 (7.4%) were positive for influenza.
- O Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the epidemic threshold.
- O <u>Influenza-associated Pediatric Deaths:</u> No influenza-associated pediatric deaths were reported.
- Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) was 1.6%, which is below the national baseline of 2.0%. All 10 regions reported ILI below region-specific baseline levels. Puerto Rico experienced moderate ILI activity; two states experienced low ILI activity; New York City and 48 states experienced minimal ILI activity; and the District of Columbia had insufficient data.
- Geographic Spread of Influenza: The geographic spread of influenza in Guam was reported as widespread; three states reported regional activity; Puerto Rico, the U.S. Virgin Islands, and 13 states reported local activity; the District of Columbia and 31 states reported sporadic activity; and three states reported no influenza activity.



